arXiv:1702.00961v1 [math.MG] 3 Feb 2017

LIOUVILLE THEOREM FOR BOUNDED HARMONIC FUNCTIONS ON GRAPHS SATISFYING NON-NEGATIVE CURVATURE DIMENSION CONDITION

BOBO HUA

Abstract. In a general setting, we prove that any bounded harmonic function on a graph satisfying the curvature dimension condition $\mathrm{CD}(0,\infty)$ is constant.

1. Introduction

Yau [Yau75] proved that positive harmonic functions are constant on a complete, noncompact, connected Riemannian manifold with non-negative Ricci curvature. As a corollary, any bounded harmonic function is constant. These are called Liouville theorems for harmonic functions, regarded as generalizations of classical Liouville's theorem for bounded holomorphic functions on the complex plane. Later, Cheng and Yau [CY75] obtained a quantitative result, so-called Cheng-Yau gradient estimate for positive harmonic functions. This type of gradient estimate was generalized to positive solutions of heat equations by Li and Yau [LY86]. The space of positive (bounded resp.) harmonic functions corresponds to the Martin (Poisson resp.) boundary at infinity for random walks in the probability theory, see e.g. [Kai96, Woe09]. As a result, the Liouville theorem is equivalent to the triviality of the corresponding boundary at infinity. Liouville theorems for harmonic functions have received much attention in the literature, to cite a few [KV83, Sul83, And83, LT87, Lyo87, Gri90, Ben91, Gri91, SC92, Wan02, Ers04, Li12, Bri13]. In this paper, we study bounded harmonic functions on graphs and prove the Liouville theorem under the curvature dimension condition.

We first recall some facts on Riemannian manifolds. Let (M,g) be a d-dimensional complete Riemannian manifold without boundary with the Riemannian metric g, Δ the Laplace-Beltrami operator on M and $C_0^\infty(M)$ the space of smooth compactly supported functions on M. As is well-known, the Ricci curvature tensor of a manifold is bounded below by Kg and the dimension $d \le n$, $K \in \mathbb{R}$ and $n \in (0,\infty]$, if and only if the following Bochner inequality holds,

$$\frac{1}{2}\Delta|\nabla f|^2 \ge \frac{1}{n}(\Delta f)^2 + \langle \nabla f, \nabla \Delta f\rangle + K|\nabla f|^2, \quad \forall\ f \in C_0^\infty(M),$$

where $\nabla\cdot$ is the Riemannian gradient of a function. Bakry and Émery [BE85, Bak87] adopted this inequality as a definition of curvature notion for a general Markov semigroup, called the curvature dimension condition and denoted by $\mathrm{CD}(K,n)$, and derived various interesting analytic properties of the semigroup from this condition. For the sake of simplicity, we consider the manifold case and denote by $P_t = e^{t\Delta}$ the heat semigroup on the manifold. One can show that the $\mathrm{CD}(0,\infty)$ condition is equivalent to

$$|\nabla P_t f|^2 \le P_t |\nabla f|^2, \quad \forall\ f \in C_0^\infty(M), t > 0,$$

or

$$2t|\nabla P_t f|^2 \le P_t(f^2) - (P_t f)^2, \quad \forall\ f \in C_0^\infty(M), t > 0, \tag{1}$$

where the latter is called the reverse Poincaré inequality.

The observation of the paper is that the reverse Poincaré inequality (1) is sufficient to prove the Liouville theorem for bounded harmonic functions. The **heuristic argument** is given as follows: Let f be a bounded harmonic function on M. We may apply (1) to f, which can be justified by proper approximation. By the semigroup property and the boundedness of f, the right hand side of (1) is bounded above by a constant C independent of t. Moreover, by the harmonicity of f, $P_t f = f$, which implies that

$$|\nabla f|^2 = |\nabla P_t f|^2 \le \frac{C}{2t} \to 0, \quad t \to \infty. \tag{2}$$

This yields that $|\nabla f| \equiv 0$ and hence f is constant.

In the following, we introduce the setting of graphs, see [HL17] for notation, and apply the previous argument to the discrete case. Let (V,E) be a connected, undirected, combinatorial graph with the set of vertices V and the set of edges E. We say $x, y \in V$ are neighbors, denoted by $x \sim y$, if there is an edge connecting x and y. The graph is called *locally finite* if each vertex has finitely many neighbors. In this paper, all graphs we consider are locally finite. We assign a weight m to each vertex, $m : V \to (0,\infty)$, and a weight μ to each edge,

$$\mu : E \to (0,\infty),\ E \ni \{x,y\} \mapsto \mu_{xy} = \mu_{yx},$$

and refer to the quadruple $G = (V,E,m,\mu)$ as a *weighted graph*. We denote by $C_0(V)$ the set of finitely supported functions on V, by $\ell^p(V,m)$, $p \in [1,\infty]$, the ℓ^p spaces of functions on V with respect to the measure m, and by $\|\cdot\|_{\ell^p(V,m)}$ the ℓ^p norm of a function.

For any weighted graph $G = (V,E,m,\mu)$, we define the *Laplacian* Δ as

$$\Delta f(x) = \frac{1}{m(x)} \sum_{y \in V : y \sim x} \mu_{xy}(f(y) - f(x)), \quad \forall\ f : V \to \mathbb{R}.$$

It is known, see Keller and Lenz [KL12], that this defines a bounded linear operator on $\ell^2(V,m)$ if and only if

$$\sup_{x\in V}\frac{1}{m(x)}\sum_{y\in V:y\sim x}\mu_{xy}<\infty. \tag{3}$$

From another point of view, for a weighted graph G, it associates with a Dirichlet form with respect to the Hilbert space $\ell^2(V,m)$,

$$\begin{aligned} Q:\quad & D(Q)\times D(Q)\to\mathbb{R}\\ & (f,g)\mapsto\frac{1}{2}\sum_{x,y\in V:x\sim y}\mu_{xy}(f(y)-f(x))(g(y)-g(x)), \end{aligned}$$

where the form domain $D(Q)$ is defined as the completion of $C_0(V)$ under the norm $\|\cdot\|_Q$ given by

$$\|f\|_Q^2=\|f\|^2_{\ell^2(V,m)}+\frac{1}{2}\sum_{x,y\in V:x\sim y}\mu_{xy}(f(y)-f(x))^2,\quad \forall f\in C_0(V),$$

see [KL12]. We denote by $\mathcal{L}$ the infinitesimal generator of the Dirichlet form Q and by $P_t=e^{t\mathcal{L}}$ the corresponding C_0-semigroup. For a locally finite graph, the generator $\mathcal{L}$ coincides with the Laplacian Δ on the domain of generator $D(\mathcal{L})$ which contains $C_0(V)$, i.e. $\mathcal{L}f=\Delta f$, for any $f\in D(\mathcal{L})$, see [KL12, Theorem 6 and 9].

Note that the weights μ and m determine the properties of the Laplacian and the semigroup. Given the edge weights μ, typical choices of m are of particular interest:

- $m(x)=\sum_{y:y\sim x}\mu_{xy}$ for any $x\in V$ and the associated Laplacian is called the normalized Laplacian.
- $m\equiv 1$ and the Laplacian is called the physical (or combinatorial) Laplacian.

Now we introduce the Γ-calculus on graphs following [BGL14, LY10]. The "carré du champ" operator Γ is defined as

$$\Gamma(f,g)(x):=\frac{1}{2}(\Delta(fg)-f\Delta g-g\Delta f)(x),\quad \forall f,g:V\to\mathbb{R},x\in V.$$

For simplicity, we write $\Gamma(f):=\Gamma(f,f)$. The iterated Γ operator, Γ_2, is defined as

$$\Gamma_2(f):=\frac{1}{2}\Delta\Gamma(f)-\Gamma(f,\Delta f).$$

Motivated by the Riemannian setting, we introduce a condition for the completeness of infinite weighted graphs: A weighted graph $G=(V,E,m,\mu)$ is called *complete* if there exists a non-decreasing sequence of finitely supported non-negative functions $\{\eta_k\}_{k=1}^{\infty}$ such that

$$\lim_{k\to\infty}\eta_k=\mathbb{1}\text{ and }\ \Gamma(\eta_k)\le\frac{1}{k}, \tag{4}$$

where $\mathbb{1}$ is the constant function 1 on V. Note that the weights μ and m appear in the definition of Γ, so that they are essential to the completeness of a weighted graph. This condition was defined for Markov diffusion semigroups in [BGL14, Definition 3.3.9] and adopted to graphs in [HL17]. A large class of graphs possessing appropriate intrinsic metrics have been shown to be complete, see [HL17, Theorem 2.8], where η_k, mimicking cut-off functions in Riemannian manifolds, are constructed via distance functions to fixed vertices. In particular, graphs with bounded Laplacians are always complete.

In this paper, we say that a weighted graph G satisfies the assumption (A) if one of the following holds:

- (A_1) The Laplacian Δ is bounded on $\ell^2(V,m)$, i.e. (3) holds.
- (A_2) G is complete and m is non-degenerate, i.e. $\inf_{x\in V} m(x) > 0$.

Note that normalized Laplacians are always bounded, hence satisfy (A_1); physical Laplacians satisfy (A_2) provided that they are complete.

For our purposes, we need the curvature dimension condition on graphs, which was initiated in [BE85, Bak87] on manifolds (or Markov diffusion semigroups) and introduced to graphs by [LY10]. We say that a graph $G=(V,E,m,\mu)$ satisfies the $\mathrm{CD}(K,n)$ condition, for $K\in\mathbb{R}$ and $n\in(0,\infty]$, if

$$\Gamma_2(f)(x) \geq \frac{1}{n}(\Delta f)^2(x) + K\Gamma(f)(x), \quad \forall\ f: V\to\mathbb{R},\ x\in V.$$

Now we are ready to state our main result.

Theorem 1. *Let $G=(V,E,m,\mu)$ be a weighted graph satisfying the assumption (A) and the $\mathrm{CD}(0,\infty)$ condition. Then any bounded harmonic function on G is constant.*

We remark that the Liouville theorem for bounded harmonic functions on graphs can be derived from the Li-Yau gradient estimate or the Harnack inequality. Both of them have been obtained by [BHL$^+$15, HLLY14] under other curvature dimension conditions such as $\mathrm{CDE}(0,n)$ and $\mathrm{CDE}'(0,n)$ for bounded Laplacians and finite n. Our assumption (A) is needed for the following result, which was proved for finite graphs by [LP14], for graphs with bounded Laplacians by [LL15], and for graphs with unbounded Laplacians satisfying (A_2) by [HL17] and [GL15].

Theorem 2. *For a weighted graph $G=(V,E,m,\mu)$ satisfying the assumption (A), the following are equivalent:*

(a) G satisfies $\mathrm{CD}(0,\infty)$.
(b) $\Gamma(P_tf) \leq P_t(\Gamma(f)), \qquad \forall\ f\in C_0(V), t>0.$
(c) $P_t(f^2)-(P_tf)^2 \leq 2tP_t(\Gamma(f)), \quad \forall\ f\in C_0(V), t>0.$
(d) $2t\Gamma(P_tf) \leq P_t(f^2)-(P_tf)^2, \quad \forall\ f\in C_0(V), t>0.$

The general results on equivalent conditions for the $\mathrm{CD}(K,n)$ condition, $K\in\mathbb{R}$ and $n\in(0,\infty]$, can be obtained as in [Wan11, LL15], which are

omitted here. We will prove Theorem 1 using (d) in Theorem 2 following the argument in (2).

We end the introduction by proposing a folklore conjecture on Liouville theorem for positive harmonic functions.

Conjecture 3. *Let $G = (V, E, m, \mu)$ be a weighted graph satisfying the assumption (A) and the $\mathrm{CD}(0, \infty)$ condition. Then any positive harmonic function on G is constant.*

2. Proof of main theorems

In this section, we give the proofs of main results.

Proof of Theorem 2. For bounded Laplacians, the theorem has been proved in [LL15, Theorem 3.1]. It suffices to prove the result under the assumption of (A_2). The equivalence of (a) and (b) was obtained in [HL17, Theorem 1.1]. The equivalence of (a) and (c) was proved in [GL15, Theorem 3.2].

We prove $(a) \Longrightarrow (d)$ following [Wan11, Theorem 1.1] or [LL15, Theorem 3.1]. The reverse direction is omitted here since it follows verbatim. Let $f \in C_0(V)$. We claim that $(P_t f)^2 \in D(\mathcal{L})$ for any $t \geq 0$. By the assumption (A_2), [KL12, Theorem 5] yields that

$$D(\mathcal{L}) = \{u \in \ell^2(V, m) : \Delta u \in \ell^2(V, m)\}.$$

On one hand, one can see that $(P_t f)^2 \in \ell^2(V, m)$ since

$$\|P_t f\|_{\ell^p(V,m)} \leq \|f\|_{\ell^p(V,m)}, \quad \forall\ p \in [1, \infty].$$

On the other hand,

$$\Delta(P_t f)^2 = 2P_t f \Delta P_t f + 2\Gamma(P_t f).$$

Note that $\Delta P_t f \in \ell^2(V, m)$ since $P_t f \in D(\mathcal{L})$. Moreover, by [HL17, Proposition 3.7],

$$\|\Gamma(P_t f)\|_{\ell^1(V,m)} \leq \|\Gamma(f)\|_{\ell^1(V,m)} < \infty.$$

Hence $\Gamma(P_t f) \in \ell^2(V, m)$ which follows from $\ell^1(V, m) \subset \ell^2(V, m)$, derived from the assumption (A_2). Combining these facts, we have $\Delta(P_t f)^2 \in \ell^2(V, m)$ and prove the claim.

Given $t > 0$, set $\phi(s, x) = P_s(P_{t-s} f)^2(x),\ 0 \leq s \leq t$. Direct calculation shows that

$$\begin{aligned} \frac{d}{ds}\phi &= \Delta P_s(P_{t-s} f)^2 - P_s(2P_{t-s} f \Delta P_{t-s} f) \\ &= P_s\left[\Delta(P_{t-s} f)^2 - 2P_{t-s} f \Delta P_{t-s} f\right] = 2P_s\Gamma(P_{t-s} f), \end{aligned}$$

where the second equality follows from that $\Delta P_s(P_{t-s} f)^2 = P_s \Delta(P_{t-s} f)^2$ since $(P_{t-s} f)^2 \in D(\mathcal{L})$, proved in the previous claim. Using (b) in Theorem 2] with the approximation of $P_{t-s} f$ by finitely supported functions,

$$\frac{d}{ds}\phi = 2P_s\Gamma(P_{t-s} f) \geq 2\Gamma(P_s P_{t-s} f) = 2\Gamma(P_t f).$$

Integration over s on $[0, t]$ gives the result (d).

□

Lemma 4. *Let $G = (V, E, m, \mu)$ be a weighted graph satisfying the assumption (A) and the $\mathrm{CD}(0, \infty)$ condition. Then for any bounded harmonic function f, $P_t f = f$.*

Proof. By (b) in Theorem 2, the semigroup P_t is stochastically complete, see [HL17, Theorem 1.2]. So that any bounded solution to the heat equation $\partial_t u = \Delta u$ is unique, see [Hua11, Theorem 1.5.1], [KL12, Theorem 1] or [Gri99]. Note that by the harmonicity of f, $u(t, x) = f(x)$ solves the heat equation with initial data f which is bounded for all $t \geq 0$. Hence $P_t f = u = f$. □

Now we can prove the Liouville theorem for bounded harmonic functions for non-negatively curved graphs.

Proof of Theorem 1. By the heuristic argument (2) in the introduction and Lemma 4, it suffices to prove that (d) in Theorem 2 holds for any bounded function f. By the completeness of the graph, there exists cut-off functions $\{\eta_k\}_{k=1}^{\infty} \subset C_0(V)$ satisfying (4). Set $f_k := f \cdot \eta_k$, for $k \geq 1$. Note that $f_k \to f$, $k \to \infty$, pointwise. By (d) in Theorem 2,

$$2t\Gamma(P_t f_k) \leq P_t(f_k^2) - (P_t f_k)^2. \tag{5}$$

Since f is bounded, the dominated convergence theorem yields that

$$P_t((f_k)^a) \to P_t(f^a), \quad k \to \infty, \quad a = 1, 2.$$

Moreover, noting that the graph is locally finite, for any $x \in V$, we have

$$\Gamma(P_t(f_k))(x) \to \Gamma(P_t f)(x), \quad k \to \infty,$$

since both sides involve only finitely many summands. Passing to the limit, $k \to \infty$, in (5), we prove the result and hence the theorem. □

Acknowledgements. B. H. is supported by NSFC, grant no. 11401106.

References

[And83] M. Anderson. The Dirichlet problem at infinity for manifolds of negative curvature. *J. Differential Geom.*, 18(4):701–721, 1983.

[Bak87] D. Bakry. Étude des transformations de Riesz dans les variétés Riemanniennes à courbure de Ricci minorée. In *Séminaire de Probabilités, XXI*, volume 1247 of *Lecture Notes in Math.*, pages 137–172, Berlin, 1987. Springer.

[BE85] D. Bakry and M. Émery. Diffusions hypercontractives. In *Séminaire de Probabilités, XIX (1983/84)*, volume 1123 of *Lecture Notes in Math.*, pages 177–206, Berlin, 1985. Springer.

[Ben91] I. Benjamini. Instability of the Liouville property for quasi-isometric graphs and manifolds of polynomial volume growth. *J. Theoret. Probab.*, 4(3):631–637, 1991.

[BGL14] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and geometry of Markov diffusion operators*. Number 348 in Grundlehren der Mathematischen Wissenschaften. Springer, Cham, 2014.

[BHL+15] F. Bauer, P. Horn, Y. Lin, G. Lippner, D. Mangoubi, and S. T. Yau. Li-Yau inequality on graphs. *J. Differential Geom.*, 99(3):359–405, 2015.

[Bri13] K. Brighton. A Liouville-type theorem for smooth metric measure spaces. *J. Geom. Anal.*, 23(2):562–570, 2013.

[CY75] S. Y. Cheng and S. T. Yau. Differential equations on Riemannian manifolds and their geometric applications. *Comm. Pure Appl. Math.*, 28(3):333–354, 1975.

[Ers04] A. Erschler. Liouville property for groups and manifolds. *Invent. Math.*, 155(1):55–80, 2004.

[GL15] C. Gong and Y. Lin. Properties for CD inequalities with unbounded Laplacians. *arXiv:1512.02471*, 2015.

[Gri90] A. Grigor'yan. Dimension of spaces of harmonic functions. *Math. Notes*, 48:1114–1118, 1990.

[Gri91] A. Grigor'yan. The heat equation on noncompact Riemannian manifolds (Russian). *Mat. Sb.*, 182(1):55–87 (English translation in Math. USSR–Sb. 72(1): 47–77, 1992.), 1991.

[Gri99] A. Grigor'yan. Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds. *Bull. Amer. Math. Soc. (N.S.)*, 36(2):135–249, 1999.

[HL17] B. Hua and Y. Lin. Stochastic completeness for graphs with curvature dimension conditions. *Adv. Math.*, 306:279–302, 2017.

[HLLY14] P. Horn, Y. Lin, S. Liu, and S. T. Yau. Volume doubling, Poincaré inequality and Guassian heat kernel estimate for nonnegative curvature graphs. *arXiv:1411.5087*, 2014.

[Hua11] X. Huang. On stochastic completeness of weighted graphs. *PhD thesis, Bielefeld University*, 2011.

[Kai96] V. A. Kaimanovich. Boundaries of invariant Markov operators: the identification problem. In M. Pollicott and K. Schmidt, editors, *Ergodic theory of $\mathbb{Z}^d$ actions (Warwick, 1993–1994)*, London Math. Soc. Lecture Note Ser. 228., pages 127–176, Cambridge, 1996. Cambridge Univ. Press.

[KL12] M. Keller and D. Lenz. Dirichlet forms and stochastic completeness of graphs and subgraphs. *J. Reine Angew. Math.*, 666:189–223, 2012.

[KV83] V. A. Kaimanovich and A. M. Vershik. Random walks on discrete groups: boundary and Entropy. *Ann. Probab.*, 11(3):457–490, 1983.

[Li12] P. Li. *Geometric analysis*, volume 134 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2012.

[LL15] Y. Lin and S. Liu. Equivalent properties of CD inequality on graph. *arXiv:1512.02677*, 2015.

[LP14] S. P. Liu and N. Peyerimhoff. Eigenvalue ratios of nonnegatively curved graphs. *arXiv:1406.6617*, 2014.

[LT87] P. Li and L. F. Tam. Positive harmonic functions on complete manifolds with nonnegative curvature outside a compact set. *Ann. of Math. (2)*, 125(1):171–207, 1987.

[LY86] P. Li and S. T. Yau. On the parabolic kernel of the Schroedinger operator. *Acta Math.*, 156(3–4):153–201, 1986.

[LY10] Y. Lin and S. T. Yau. Ricci curvature and eigenvalue estimate on locally finite graphs. *Math. Res. Lett.*, 17(2):343–356, 2010.

[Lyo87] T. Lyons. Instability of the Liouville property for quasi-isometric Riemannian manifolds and reversible Markov chains. *J. Differential Geom.*, 26(1):33–66, 1987.

[SC92] L. Saloff-Coste. A note on Poincaré, Sobolev, and Harnack inequalities. *Internat. Math. Res. Notices*, (2):27–38, 1992.

[Sul83] D. Sullivan. The Dirichlet problem at infinity for a negatively curved manifold. *J. Differential Geom.*, 18(4):723–732, 1983.

[Wan02] F. Y. Wang. Liouville theorem and coupling on negatively curved manifolds. *Stochastic Process. Appl.*, 100:27–39, 2002.

[Wan11] F. Y. Wang. Equivalent semigroup properties for curvature-dimension condition. *Bull. Sci. Math.*, 135(6-7):803–815, 2011.

[Woe09] W. Woess. *Denumerable Markov chains–Generating functions, boundary theory, random walks on trees.* EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2009.

[Yau75] S. T. Yau. Harmonic functions on complete Riemannian manifolds. *Comm. Pure Appl. Math.*, 28:201–228, 1975.

E-mail address: bobohua@fudan.edu.cn

School of Mathematical Sciences, LMNS, Fudan University, Shanghai 200433, China; Shanghai Center for Mathematical Sciences, Fudan University, Shanghai 200433, China